

Effective refractive index tensor for weak-field gravity

Petarpa Boonserm [†], Celine Cattoen [‡], Tristan Faber [§],
Matt Visser ^{||}, and Silke Weinfurtner [¶]

School of Mathematics, Statistics, and Computer Science,
Victoria University of Wellington,
P.O.Box 600, Wellington, New Zealand

Abstract.

Gravitational lensing in a weak but otherwise arbitrary gravitational field can be described in terms of a 3×3 tensor, the “effective refractive index”. If the sources generating the gravitational field all have small internal fluxes, stresses, and pressures, then this tensor is automatically isotropic and the “effective refractive index” is simply a scalar that can be determined in terms of a classic result involving the Newtonian gravitational potential. In contrast if anisotropic stresses are ever important then the gravitational field acts similarly to an anisotropic crystal. We derive simple formulae for the refractive index tensor, and indicate some situations in which this will be important.

gr-qc/0411034; 8 November 2004;

Revised 10 March 2005; L^AT_EX-ed 7 February 2008

PACS numbers: 04.20.-q; 04.20.Cv

[†] petarpa.boonserm@mcs.vuw.ac.nz

[‡] celine.cattoen@mcs.vuw.ac.nz

[§] tristan.faber@mcs.vuw.ac.nz

^{||} matt.visser@mcs.vuw.ac.nz

[¶] silke.weinfurtner@mcs.vuw.ac.nz

1. Introduction

Weak-field gravity in Einstein’s general relativity is actually more general than straightforward Newtonian gravity [1, 2]. While the approximate validity of Newtonian gravity is certainly limited to the weak-field regime, Newtonian gravity makes significant additional assumptions as to the smallness of effects that depend on the internal stresses, pressures, and energy fluxes in the massive bodies that act as source for the gravitational field. While there is no significant doubt that for planets, and most stars, the gravitational effects of internal stresses can safely be neglected, the situation for neutron stars (where $G_N M/R \approx 1/10$) is much more uncertain. Furthermore, while there is little doubt that the “dark matter” that makes up approximately 90% of most spiral galaxies can be treated using weak-field gravity, in the absence of solid physical motivation for some particular equation of state we cannot necessarily conclude that the gravitational field can be adequately described by Newtonian gravity.

In view of this we have developed a formalism that makes no assumptions about the relative smallness or isotropy of internal stresses (and pressures and fluxes), to see how gravitational lensing is affected. In particular, weak Newtonian gravitational lenses can be interpreted in terms of an analogy wherein a gravitational field is assigned an “effective refractive index” [3, 4], and we extend these ideas to see how this “effective refractive index” is affected by the presence of significant internal stress. Most strikingly we will see that the “effective refractive index” is in general no longer a scalar, but is instead a 3×3 tensor — in analogy to the situation in an anisotropic crystal. (The use of analogies to relate otherwise distinct phenomena, and to give qualitative insight as to what physical effects might be important, has recently attracted significant interest in the general relativity community [5], but related ideas under the name “electro-optical analogy” have an independent history [6].) We organize the paper as follows:

- First we consider the static case where there are no internal energy fluxes (so in particular we neglect the effects of rotation).
- Second we further specialise this discussion to situations of static spherical symmetry — believed to be a good approximation for galactic halos containing “dark matter”.
- Third we extend the discussion to the more general stationary case, where internal fluxes are included.
- Fourth we indicate how time-dependent situations can in principle be dealt with.
- Fifth we indicate how the present formalism matches to the usual idea of a far-field multipole expansion.
- Finally we briefly discuss astrophysical situations in which the issues raised in this article are likely to become important.

2. Static case

For light propagating in curved space along some curve parameterized by λ we have

$$g_{ab} dX^a(\lambda) dX^b(\lambda) = 0. \quad (1)$$

Looking at the specific case of a weak field, where $g_{ab} = \eta_{ab} + h_{ab}$, the gravitational field can be considered as a perturbation h_{ab} around the flat space η_{ab} . This leads to:

$$g_{ab} dX^a(\lambda) dX^b(\lambda) = \eta_{ab} dX^a(\lambda) dX^b(\lambda) + h_{ab} dX^a(\lambda) dX^b(\lambda) = 0. \quad (2)$$

Here $\eta_{ab} dX^a(\lambda) dX^b(\lambda)$ is no longer zero. For a light ray propagating in a *static* weak field we get

$$g_{ab} \frac{dX^a}{d\lambda} \frac{dX^b}{d\lambda} = (-1 + h_{tt}) \left(\frac{dt}{d\lambda} \right)^2 + (\delta_{ij} + h_{ij}) \frac{dx^i}{d\lambda} \frac{dx^j}{d\lambda} = 0. \quad (3)$$

Choosing $\lambda = t$ equation (3) simplifies to

$$g_{ab} \frac{dX^a}{dt} \frac{dX^b}{dt} = (-1 + h_{tt}) + (\delta_{ij} + h_{ij}) \dot{x}^i \dot{x}^j = 0. \quad (4)$$

We define a “coordinate speed of light” by calculating the norm $||\dot{x}^i||$ of the “coordinate velocity of light” $\dot{x}^i$ using δ_{ij} , the unperturbed background metric for space. This allows us to split the velocity into a speed and a direction

$$\dot{x}^i = ||\dot{x}^i|| \hat{k}^i; \quad ||\hat{k}^i|| = 1 = \sqrt{\delta_{ij} \hat{k}^i \hat{k}^j}; \quad (5)$$

where $\hat{k}^i$ is a unit 3-vector. Putting this into equation (4), and noting that the h_{ab} are small compared to unity we can usefully Taylor series expand, to obtain

$$||\dot{x}^i|| = \sqrt{\frac{1 - h_{tt}}{1 + h_{ij} \hat{k}^i \hat{k}^j}} \approx 1 - \frac{1}{2} h_{tt} - \frac{1}{2} h_{ij} \hat{k}^i \hat{k}^j + \mathcal{O}(h_{ab}^2) \quad (6)$$

for the coordinate speed of light. Note that we have adopted units where the physical speed of light, c , measured by physical rulers and physical clocks, is always 1. Then the spacetime refractive index for light travelling in the direction $\hat{k}$ is

$$n(\hat{k}) = \frac{1}{||\dot{x}^i||} \approx 1 + \frac{1}{2} h_{tt} + \frac{1}{2} h_{ij} \hat{k}^i \hat{k}^j + \mathcal{O}(h_{ab}^2). \quad (7)$$

We now define the 3×3 *refractive index tensor* as:

$$n_{ij} \equiv \left(1 + \frac{1}{2} h_{tt} \right) \delta_{ij} + \frac{1}{2} h_{ij}, \quad (8)$$

so that $n(\hat{k}) = n_{ij} \hat{k}^i \hat{k}^j + \mathcal{O}(h_{ab}^2)$. If we adopt the standard definition

$$\bar{h}^{ab} = h^{ab} - \frac{1}{2} h \eta^{ab}, \quad (9)$$

then a brief computation yields

$$n_{ij} \equiv \left(1 + \frac{1}{2} \bar{h}_{tt} \right) \delta_{ij} + \frac{1}{2} \bar{h}_{ij}. \quad (10)$$

To connect these general formulae to the presence of stress-energy, adopt Einstein–Fock–de Donder gauge (that is, harmonic quasi-Cartesian coordinates) [1]

$$\partial_a \bar{h}^{ab} = 0, \quad (11)$$

and write the Einstein equations in the *exact* form [1]

$$\nabla^2 \bar{h}_{ab} = -16\pi G_N T_{ab}^{\text{eff}}, \quad (12)$$

where the “effective” stress-energy contains contributions both from “ordinary” stress-energy and the “pseudo-energy” of the gravitational field itself:

$$T_{ab}^{\text{eff}} = T_{ab} + t_{ab}. \quad (13)$$

In all situations we are interested in the pseudo-energy is much smaller than the ordinary stress energy, nevertheless explicitly keeping the pseudo-energy as part of the effective stress energy is a very useful bookkeeping device. (This is an application of the “principle of controlled ignorance” espoused in [1].) Due to the assumption that the spacetime is static, the effective stress-energy tensor is

$$T_{ab}^{\text{eff}} = \begin{bmatrix} \rho^{\text{eff}} & 0 \\ 0 & T_{ij}^{\text{eff}} \end{bmatrix}. \quad (14)$$

We define

$$\nabla^2 \Phi \equiv 4\pi G_N \rho^{\text{eff}}, \quad (15)$$

$$\nabla^2 \Psi_{ij} \equiv 4\pi G_N T_{ij}^{\text{eff}}, \quad (16)$$

where Φ is the ordinary Newton potential, but with the effective mass-energy-density ρ^{eff} as a source. The Ψ_{ij} are new post-Newton gravitational potentials arising from the effective internal pressures and stresses of the source matter. Using the Einstein equations we find:

$$\nabla^2 \bar{h}_{ij} = -16\pi G_N T_{ij}^{\text{eff}}. \quad (17)$$

$$\nabla^2 \bar{h}_{tt} = -16\pi G_N \rho^{\text{eff}}. \quad (18)$$

Imposing suitable boundary conditions at spatial infinity and working in terms of the density and pressure potentials, we obtain

$$\bar{h}_{tt} = -4\Phi; \quad (19)$$

$$\bar{h}_{ij} = -4\Psi_{ij}, \quad (20)$$

and the equivalent (though more complicated)

$$h_{tt} = -2(\Phi + \delta^{kl} \Psi_{kl}); \quad (21)$$

$$h_{ij} = -2(2\Psi_{ij} + \delta_{ij}[\Phi - \delta^{kl} \Psi_{kl}]). \quad (22)$$

Thus, the refractive index tensor defined in (8,10) takes the particularly simple form

$$n_{ij} = (1 - 2\Phi) \delta_{ij} - 2\Psi_{ij}. \quad (23)$$

In view of our use of the effective stress-energy, these statements are now all exact, but formal. Because h_{ab} and its derivatives occur in the pseudo-tensor t_{ab} , the RHS of

these formulae depend implicitly on h_{ab} . In practical calculations one would start by setting $h_{ab} \rightarrow 0$ on the RHS, and then iterate the equations to obtain the desired level of accuracy.

For instance, if the effective internal stresses are isotropic $T_{ij}^{\text{eff}} \rightarrow p^{\text{eff}} \delta_{ij}$ and $\Psi_{ij} \rightarrow \Psi_0 \delta_{ij}$, we have the simplification

$$h_{tt} = -2(\Phi + 3\Psi_0); \quad (24)$$

$$h_{ij} = -2\delta_{ij}[\Phi - \Psi_0]; \quad (25)$$

$$n_{ij} = (1 - 2\Phi - 2\Psi_0) \delta_{ij}, \quad (26)$$

with

$$\nabla^2 \Phi = 4\pi G_N \rho^{\text{eff}}, \quad (27)$$

$$\nabla^2 \Psi_0 = 4\pi G_N p^{\text{eff}}. \quad (28)$$

Thus for isotropic effective stress, the refractive index is a scalar. If furthermore the gravitational field is weak and smooth enough that $t_{ab} \ll T_{ab}$ this can be turned into an approximate statement about physical perfect fluids, where $T_{ij} \rightarrow p \delta_{ij}$ — the refractive index would now be approximately isotropic. (This is the usual situation in most stars: the stellar material is to a good approximation a perfect fluid and the gravitational pseudo-energy is a small fraction of the total mass budget.)

Finally, if the internal stresses are completely negligible (which is the usual situation in most asteroids, but not stars and planets [1]) we can naively set $\Psi_{ij} \rightarrow 0$ to obtain the Newtonian limit and recover the well-known standard results [3, 4]:

$$h_{tt} = -2\Phi; \quad h_{ij} = -2\Phi \delta_{ij}; \quad n_{ij} = (1 - 2\Phi) \delta_{ij}. \quad (29)$$

The novelty in the current analysis lies exactly in the manner in which internal stresses in the body generating the gravitational field lead to a “stress potential” Ψ_{ij} which then influences both the weak-field metric (19–22) and the effective refractive index tensor (23).

3. Static spherically symmetric weak field

These general considerations can be made more explicit by working in situations of spherical symmetry. Consider now a general static spherically symmetric weak field, for which the stress energy tensor (with the background Minkowski metric η_{ab} written in terms of spherical polar coordinates) takes the form

$$T_{ab}^{\text{eff}} = \begin{bmatrix} \rho^{\text{eff}}(r) & 0 & 0 & 0 \\ 0 & p_r^{\text{eff}}(r) & 0 & 0 \\ 0 & 0 & p_t^{\text{eff}}(r) r^2 & 0 \\ 0 & 0 & 0 & p_t^{\text{eff}}(r) r^2 \sin^2 \theta \end{bmatrix}. \quad (30)$$

Here p_r^{eff} and p_t^{eff} are the radial and transverse pressures respectively. At the origin we must have $p_r^{\text{eff}}(0) = p_t^{\text{eff}}(0)$. Because of the way the Einstein equations interact with the harmonic gauge condition, T_{eff}^{ab} is exactly conserved in the flat space covariant sense:

$$[T_{\text{eff}}^{ab}]_{;b} = 0; \quad \text{implying} \quad T^{ab}_{;b} = -t^{ab}_{;b} \quad (31)$$

where the colon denotes the flat-space covariant derivative corresponding to η_{ab} in spherical polar coordinates. Consequently

$$\partial_r p_r^{\text{eff}}(r) + \frac{2\{p_r^{\text{eff}}(r) - p_t^{\text{eff}}(r)\}}{r} = 0, \quad (32)$$

which we can use to eliminate $p_t(r)$ as

$$p_t^{\text{eff}}(r) = p_r^{\text{eff}}(r) + \frac{1}{2} r \partial_r p_r^{\text{eff}}(r). \quad (33)$$

We can similarly write

$$\bar{h}_{ab} = \begin{bmatrix} H_0(r) & 0 & 0 & 0 \\ 0 & H_r(r) & 0 & 0 \\ 0 & 0 & H_t(r)r^2 & 0 \\ 0 & 0 & 0 & H_t(r)r^2 \sin^2 \theta \end{bmatrix}, \quad (34)$$

where at the origin symmetry demands $H_r(0) = H_t(0)$, and the harmonic gauge condition (11), now in the flat-space spherical polar sense $\bar{h}^{ab}{}_{;b} = 0$, implies

$$H_t(r) = H_r(r) + \frac{1}{2} r \partial_r H_r(r). \quad (35)$$

After a brief computation

$$\nabla^2 \bar{h}_{ab} = \bar{h}_{ab;c}{}^{;c} = \begin{bmatrix} Z_0(r) & 0 & 0 & 0 \\ 0 & Z_r(r) & 0 & 0 \\ 0 & 0 & Z_t(r)r^2 & 0 \\ 0 & 0 & 0 & Z_t(r)r^2 \sin^2 \theta \end{bmatrix}, \quad (36)$$

where we are again using the covariant derivative for flat-space spherical polar coordinates, and we have defined

$$Z_0 = \frac{1}{r^2} \partial_r (r^2 \partial_r H_0); \quad (37)$$

$$Z_r = \frac{1}{r^2} \partial_r (r^2 \partial_r H_r) - 4 \frac{H_r(r) - H_t(r)}{r^2} = \frac{1}{r^4} \partial_r (r^4 \partial_r H_r); \quad (38)$$

$$Z_t = \frac{1}{r^2} \partial_r (r^2 \partial_r H_t) + 2 \frac{H_r(r) - H_t(r)}{r^2} = Z_r + \frac{1}{2} r \partial_r Z_r. \quad (39)$$

The slightly unusual terms proportional to the difference $H_r - H_t$ arise due to the fact that we are now using spherical polar coordinates (not strictly harmonic coordinates), and because h_{ab} has tensor indices in its own right. The Einstein equations now yield

$$Z_0 = -16\pi G_N \rho^{\text{eff}}; \quad (40)$$

$$Z_r = -16\pi G_N p_r^{\text{eff}}; \quad (41)$$

plus the redundant equation

$$Z_t = -16\pi G_N p_t^{\text{eff}}. \quad (42)$$

These have the formal solutions

$$H_0 = -16\pi G_N \int \left\{ \int \rho^{\text{eff}} r^2 dr \right\} r^{-2} dr = -4\Phi; \quad (43)$$

$$H_r = -16\pi G_N \int \left\{ \int p_r^{\text{eff}} r^4 dr \right\} r^{-4} dr; \quad (44)$$

where we have again used equation (15), and so

$$H_r - H_t = \frac{8\pi G_N}{r^3} \left\{ \int p_r^{\text{eff}} r^4 dr \right\}. \quad (45)$$

We can further rearrange this result slightly by defining

$$H_\Sigma \equiv H_r + 2H_t; \quad H_\Delta \equiv H_r - H_t, \quad (46)$$

so that

$$H_r = \frac{H_\Sigma + 2H_\Delta}{3}; \quad H_t = \frac{H_\Sigma - H_\Delta}{3}. \quad (47)$$

For a null curve located at radius r , and making an angle χ with respect to the $\hat{r}$ direction, equation (10) implies

$$n(r, \chi) = 1 + \frac{1}{2}H_0 + \frac{1}{2}(H_r \cos^2 \chi + H_t \sin^2 \chi), \quad (48)$$

so that

$$n(r, \chi) = 1 + \frac{1}{2}H_0 + \frac{1}{6}H_\Sigma + \frac{1}{6}H_\Delta \{3 \cos^2 \chi - 1\}, \quad (49)$$

or

$$n(r, \chi) = 1 - 2\Phi + \frac{1}{6}H_\Sigma + \frac{1}{6}H_\Delta \{3 \cos^2 \chi - 1\}. \quad (50)$$

This particularises equation (23) to the case of spherical symmetry, with Ψ_{ij} now being written in terms of H_Σ and H_Δ , and with explicit integral formulae now being available for these quantities.

Let's compare this with a situation where we know the exact result: If the object that acts as source for the gravitational field has a definite surface, with vacuum outside that surface, then application of the Birkhoff theorem tells us that the spacetime geometry will be Schwarzschild outside that surface. But because of our subsidiary assumption, that we are working in the spherical polar version of Einstein–Fock–de Donder gauge, there is no remaining freedom in our coordinate system and we must obtain the weak-field limit of Schwarzschild geometry in harmonic coordinates — for which the effective refractive index is asymptotically isotropic. In harmonic coordinates the Schwarzschild solution has

$$h_{rr} = \frac{2m}{r} - \frac{2m^2}{r^2} + \mathcal{O}(r^{-3}); \quad (51)$$

$$\frac{h_{\theta\theta}}{r^2} = \frac{h_{\phi\phi}}{r^2 \sin^2 \theta} = \frac{2m}{r} + \frac{m^2}{r^2}; \quad (52)$$

$$H_\Delta = -\frac{3m^2}{r^2} + \mathcal{O}(r^{-3}). \quad (53)$$

Hence anisotropies in the effective refractive index are (at least in the case of spherical symmetry) constrained to rapidly decay outside the source that generates them. We now show that this is compatible with equation (45) above. Outside the body $T_{ab} = 0$ so

$p_r^{\text{eff}} \rightarrow t_{rr}$ depends on the pseudo-stress only. But $t_{rr} \sim (\partial h)^2 \sim (m/r^2)^2 = m^2/r^4$. This then implies $H_\Delta \sim m^2/r^2$, in agreement with the exact result. The precise distribution of effective refractive index anisotropies inside the source object can only be determined by solving the Einstein equations, which requires some specific model for the interior distribution of anisotropic stresses. This is a topic which we hope to explore more fully in the future.

4. Stationary case

Let us now consider the stationary case, where

$$h_{ab} = \begin{bmatrix} h_{tt} & h_{tj} \\ h_{it} & h_{ij} \end{bmatrix}, \quad (54)$$

and due to symmetry $h_{tj} = h_{jt}$. The condition for a photon trajectory then becomes:

$$g_{ab} \frac{dX^a}{d\lambda} \frac{dX^b}{d\lambda} = (-1 + h_{tt}) \left(\frac{dt}{d\lambda} \right)^2 + 2h_{tj} \frac{dt}{d\lambda} \frac{dx^j}{d\lambda} + (\delta_{ij} + h_{ij}) \frac{dx^i}{d\lambda} \frac{dx^j}{d\lambda} = 0. \quad (55)$$

Choosing the parameter $\lambda = t$, this becomes a quadratic equation in $\dot{x}^i$ with the coordinate speed of light now being given by

$$||\dot{x}^i|| = \frac{1}{(\delta_{ij} + h_{ij}) \hat{k}^i \hat{k}^j} \left(-h_{tj} \hat{k}^j + \sqrt{(h_{tj} \hat{k}^j)^2 - (-1 + h_{tt})(\delta_{ij} + h_{ij}) \hat{k}^i \hat{k}^j} \right). \quad (56)$$

Simplifying, Taylor expanding, and inverting gives the refractive index

$$n(\hat{k}) = 1 + \frac{1}{2} \left(h_{tt} + 2h_{tj} \hat{k}^j + h_{ij} \hat{k}^i \hat{k}^j \right) + \mathcal{O}(h^2), \quad (57)$$

which is a very straightforward extension of the static case. But because of the linear term in $\hat{k}$, it is not possible to bring this completely into 3-tensor form — there is additional structure and we must write

$$n(\hat{k}) = n_{ij} \hat{k}^i \hat{k}^j + h_{tj} \hat{k}^j + \mathcal{O}(h^2), \quad (58)$$

where n_{ij} has exactly the same form as in the static case [see equation (8)] and the new $h_{tj} \hat{k}^j$ term can be interpreted as being due to motion of the “effective medium” with respect to the quasi-Cartesian coordinate system (t, x^i) . This can be justified by performing a coordinate transformation into the local rest frame of the “effective medium”, which is moving with “velocity” $-h_{tj}$, and exhibits a refractive index tensor n_{ij} . See Appendix A for details. Since the “medium” is generally moving inhomogeneously the local rest frame is not best for performing explicit calculations — for practical calculations it is preferable to fix the coordinate system once and for all and to work with the $n(\hat{k})$ of equation (58) above. (In particular, the differential equations relating the weak field h_{ab} to the distribution of stress energy are defined in the original coordinate system, which was chosen to satisfy the Einstein–Fock–de Donder gauge condition, and these differential equations do not adapt nicely to the comoving point of view.)

We can furthermore define additional “flux potentials”

$$\nabla^2 \Pi_j = 4\pi G_N T_{tj}^{\text{eff}} \quad (59)$$

that couple to the momentum flux. The corresponding weak-field Einstein equations are very simple, since η_{ab} vanishes for the off-diagonal elements:

$$\nabla^2 \bar{h}_{tj} = \nabla^2 h_{tj} = -16\pi G_N T_{tj}^{\text{eff}}. \quad (60)$$

Imposing appropriate boundary conditions

$$h_{tj} = -4\Pi_j, \quad (61)$$

and the refractive index for the stationary case is

$$n(\hat{k}) = n_{ij}(\Phi, \Psi) \hat{k}^i \hat{k}^j - 4\Pi_j \hat{k}^j + \mathcal{O}(\Phi^2, \Psi^2, \Pi^2) \quad (62)$$

where $n_{ij}(\Phi, \Psi)$ is as in equation (23).

An alternative representation is to note that if we define $k^a = (1; \hat{k}^i)$ so that k^a is a null vector with respect to the background metric, then

$$n(\hat{k}) = 1 + \frac{1}{2} h_{ab} k^a k^b + \mathcal{O}(h^2) = 1 + \frac{1}{2} \bar{h}_{ab} k^a k^b + \mathcal{O}(h^2). \quad (63)$$

If we further define

$$\Phi_{ab} \equiv \left[\begin{array}{c|c} \Phi & \Pi_i \\ \hline \Pi_j & \Psi_{ij} \end{array} \right], \quad (64)$$

then

$$n(\hat{k}) = 1 - 2\Phi_{ab} k^a k^b + \mathcal{O}(\Phi_{ab}^2). \quad (65)$$

5. Time dependent situations

In the presence of time-dependent sources the only modification is that the Laplace equations for the potentials should be replaced by wave equations $\nabla^2 \rightarrow -\partial_t^2 + \nabla^2$, which has the effect of replacing the usual $1/r$ potentials by the appropriate Liénard–Wiechert potential [7]. The net result is that

$$h_{ab}(x, t) = 16\pi G_N \int d^3y \frac{\left\{ T_{ab}^{\text{eff}}(y, \tilde{t}) - \frac{1}{2} \eta_{ab} T^{\text{eff}}(y, \tilde{t}) \right\}}{|\vec{x} - \vec{y}|}, \quad (66)$$

where $\tilde{t}$ is the retarded time

$$\tilde{t} = t - |\vec{x} - \vec{y}|. \quad (67)$$

But if the so-called null energy condition [NEC] is satisfied [7], then in particular $T_{ab} k^a k^b \geq 0$, implying $h_{ab} k^a k^b \geq \mathcal{O}(h_{ab}^2)$, so that $n(\hat{k}) \geq 1 + \mathcal{O}(h_{ab}^2)$. That is, the NEC implies the effective refractive index is [to order $\mathcal{O}(h_{ab}^2)$] greater than unity, thereby guaranteeing that in this approximation the coordinate speed of light is always less than 1. This connects the discussion back to the perturbative version of “superluminal censorship” discussed in [7].

6. The far field

For isolated bodies it is possible to expand the far field in terms of multipole moments. In particular, if we go to the rest frame of the body, and if the body has time-independent internal structure, then it is a standard result that [1]

$$h_{ab} = \frac{2G_N M}{r} \delta_{ab} + \mathcal{O}(1/r^2). \quad (68)$$

Thus for a single isolated body

$$n(\hat{k}) = 1 + \frac{1}{2} h_{ab} k^a k^b + \mathcal{O}(h^2) = 1 + \frac{2G_N M}{r} + \mathcal{O}(1/r^2), \quad (69)$$

where $k^a = (1; \hat{k}^i)$ is a null vector with respect to the background metric. But as long as the total mass is positive $M > 0$ we have $n(\hat{k}) \geq 1$. That is, the positive mass theorem implies that in the far-field regime the effective refractive index is always greater than unity, thereby guaranteeing that the coordinate speed of light is always less than 1. This is a rather different perturbative version of “superluminal censorship”, more akin to the ideas discussed in [8]. Furthermore note that for an isolated body at rest the far-field refractive index is automatically isotropic — this is consistent with our calculation for situations possessing spherical symmetry, where we found that anisotropic propagation of null geodesics was confined to the region inside the body.

If there are many compact objects making up the gravitational lens, one should simply sum the $1/r$ potentials for each object. If we are considering a body that is at rest but rotating, then the overall momentum P^i is zero and the dominant term in the potential Π_i comes from the dipole term in the multipole expansion — this term is well-known to be related to the total angular momentum of the rotating body and a standard textbook result yields [1]

$$\Pi_i = \frac{G_N (\mathbf{L} \times \hat{\mathbf{r}})_i}{2r^2} + \mathcal{O}(1/r^3). \quad (70)$$

If we consistently retain all $\mathcal{O}(1/r^2)$ terms in a post-Newtonian analysis then the far field for a stationary rotating body is (see, for example, equation (19.13) of [1])

$$h_{ab} = \left[\frac{2G_N M/r - 2G_N^2 M^2/r^2}{-2G_N (\mathbf{L} \times \hat{\mathbf{r}})_i/r^2} \middle| \frac{-2G_N (\mathbf{L} \times \hat{\mathbf{r}})_j/r^2}{(2G_N M/r + 3G_N^2 M^2/2r^2)\delta_{ij}} \right] + \mathcal{O}(1/r^3) \quad (71)$$

which implies

$$n(\hat{k}) = 1 + \frac{2G_N M}{r} - \frac{G_N^2 M^2}{4r^2} - \frac{2G_N (\mathbf{L} \times \hat{\mathbf{r}}) \cdot \hat{\mathbf{k}}}{r^2} + \mathcal{O}(1/r^3). \quad (72)$$

So angular momentum certainly does leave an imprint on the far-field refractive index, but in a sub-dominant term.

Of course if the bodies in question are not isolated (one might for instance be considering a null geodesic passing through the bulk of the body), then the multipole expansion is not useful — and one should resort to the use of the general equation (62) and the linearized field equations for Φ , Π^i , and Ψ^{ij} (with $\nabla^2 \rightarrow -\partial_t^2 + \nabla^2$ for time dependent situations).

7. Discussion

Our formulation of weak field [but not necessarily Newtonian] gravitational lensing is particularly simple and gives a nice interpretation of the “effective refractive index” of the gravitational field directly in terms of Newtonian-like potentials coupled to the stress-energy tensor. Doing so results in an “effective refractive index” more general than the standard Newtonian result — anisotropic stresses are seen to lead to an anisotropic “refractive index” and energy fluxes and angular momentum are seen to lead to a “moving medium” effect.

First we should raise (and settle) the issue of gauge invariance [coordinate independence] — our results were obtained by using the simplifying properties of Einstein–Fock–de Donder gauge, and so clearly there is a sense in which the existence and value of the refractive index tensor depends critically on the use of specific coordinates. However it must be emphasised that once one has used the refractive index to calculate quantities such as magnification factors and/or the angle measured between two images on the sky, these angles and magnifications have a coordinate-invariant physical meaning independent of whatever coordinate system was used to carry out the calculation. So while the analogy that leads to the refractive index tensor is not itself coordinate invariant, the physical observables that result at the end of any specific calculation are coordinate invariants.

We now ask under what circumstances these effects might physically be important? Are there physical situations in which the gravitational field is appreciable (but still a weak-field), but with $T_{ij} \approx T_{tt}$? Perhaps the best known situation of this type arises in the core of a neutron star where $GM/R \approx 1/10$ so the gravitational field is reasonably weak, and where matter is approximately described by a “stiff” equation of state, $p = \rho c^2$. Furthermore, a radiation fluid (or neutrino fluid) satisfies $p = \rho c^2/3$, so if radiation fluid ever becomes a significant fragment of the total mass budget of any clumped system, the ideas of this article would become important. The contribution of angular momentum to microlensing, which in our formalism arises at the dipole level in the Π_i potential, has been considered in several articles [9] using somewhat different formalism. A particularly interesting effect is the possibility of anisotropy in the refractive index — this occurs once one moves away from the perfect fluid approximation for astrophysical bodies, and speculation concerning significant crustal stresses in neutron stars (and their possible effects on compactness bounds) is common [10]. Finally we point out that coherent field configurations, be they electromagnetic fields or scalar fields, generally induce anisotropic stresses comparable in magnitude to the energy density. In particular any attempt at modelling the “dark matter” in galactic halos with classical fields will lead to anisotropic stresses that, while weak, are comparable in magnitude to the energy density [11].

In summary, so long as there is continuing uncertainty over the total mass budget and relevant equations of state for the various compact objects occurring in our universe we feel that it is prudent to retain the generality of the analysis in the present article.

Appendix A. Moving medium interpretation

To see how a stationary (non-static) weak gravitational field can be viewed as a moving “effective medium”, it is useful to temporarily work with strong gravitational fields. Pick a particular point in spacetime (t_0, x_0^i) . and consider the change of coordinates

$$t \rightarrow \bar{t} = t; \quad x^i \rightarrow \bar{x}^i = x^i - v^i(t_0, x_0) [t - t_0] \quad (\text{A.1})$$

with $v^i(t_0, x_0) = g^{ik}(t_0, x_0) g_{tk}(t_0, x_0)$. Then

$$d\bar{t} = dt; \quad d\bar{x}^i = dx^i + g^{ik}(t_0, x_0) g_{tk}(t_0, x_0) dt \quad (\text{A.2})$$

and in the immediate vicinity of (t_0, x_0^i) we have

$$g_{ab} dX^a dX^b = g_{tt} dt^2 + 2g_{ti} dt dx^i + g_{ij} dx^i dx^j \quad (\text{A.3})$$

$$= [g_{tt} - (g_{ti} g^{ij} g_{jt})] d\bar{t}^2 + g_{ij} d\bar{x}^i d\bar{x}^j \quad (\text{A.4})$$

which clearly has the effect of (locally) banishing the mixed time-space parts of the metric at the cost that

$$\bar{g}_{tt} = g_{tt} - (g_{ti} g^{ij} g_{jt}); \quad \bar{g}_{ti} = 0; \quad \bar{g}_{ij} = g_{ij}. \quad (\text{A.5})$$

But because we wish to apply this in the weak field approximation $g_{ab} = \eta_{ab} + h_{ab}$ with $h_{ab} \ll 1$, this simplifies tremendously. Since $g^{ti} = h^{ti} + \mathcal{O}(h^2)$, whereas $g_{ti} = h_{ti}$ exactly, we see

$$\bar{h}_{tt} = h_{tt} + \mathcal{O}(h^2); \quad \bar{h}_{ti} = 0; \quad \bar{h}_{ij} = h_{ij}. \quad (\text{A.6})$$

That is, in weak field going to the moving coordinates defined by

$$t \rightarrow \bar{t} = t; \quad x^i \rightarrow \bar{x}^i = x^i + \delta^{ik} h_{tk}(t_0, x_0) [t - t_0] + \mathcal{O}(h^2); \quad (\text{A.7})$$

$$d\bar{t} = dt; \quad d\bar{x}^i = dx^i + \delta^{ik} h_{tk}(t_0, x_0) dt + \mathcal{O}(h^2); \quad (\text{A.8})$$

allows us (locally) to interpret $-h_{tk}$ as the “velocity” of the medium, while h_{ij} and h_{tt} , since they change by at worst $\mathcal{O}(h^2)$, lead to the same refractive index tensor as was encountered in the static case.

Acknowledgements

This research was supported by the Marsden Fund administered by the Royal Society of New Zealand.

References

- [1] C W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation*, (Freeman, San Francisco, 1972). See esp. pp. 435 ff.
- [2] B. F. Schutz, *A first course in general relativity*, (Cambridge, England, 1985). See esp. pp. 200 ff.
- [3] P. Schneider, J. Ehlers, E.E. Falco, E. E. Falco, *Gravitational Lenses* (Springer Verlag, 1993).
- [4] Silvia Mollerach and Esteban Roulet, *Gravitational Lensing and Microlensing*, (World Scientific, Singapore, 2002).
- [5] W. G. Unruh, “Dumb holes and the effects of high frequencies on black hole evaporation,” *Phys. Rev. D* **51** (1995) 2827 [arXiv:gr-qc/9409008].
M. Visser, “Acoustic propagation in fluids: An Unexpected example of Lorentzian geometry,” arXiv:gr-qc/9311028.
M. Visser, “Acoustic black holes: Horizons, ergospheres, and Hawking radiation,” *Class. Quant. Grav.* **15** (1998) 1767 [arXiv:gr-qc/9712010].
M. Novello, M. Visser and G. Volovik, “Artificial black holes,” (World Scientific, Singapore, 2002).
- [6] K. K. Nandi, Y. Z. Zhang, P. M. Alsing, J. C. Evans and A. Bhadra, “Analog of the Fizeau Effect in an Effective Optical Medium,” *Phys. Rev. D* **67** (2003) 025002 [arXiv:gr-qc/0208035].
J. C. Evans, P. M. Alsing, S. Giorgetti and K. K. Nandi, “Matter waves in a gravitational field: An index of refraction for massive particles in general relativity,” *Am. J. Phys.* **69** (2001) 1103 [arXiv:gr-qc/0107063].
- [7] M. Visser, B. Bassett and S. Liberati, “Superluminal censorship,” *Nucl. Phys. Proc. Suppl.* **88** (2000) 267 [arXiv:gr-qc/9810026].
M. Visser, B. Bassett and S. Liberati, “Perturbative superluminal censorship and the null energy condition,” published in *General Relativity and Relativistic Astrophysics, Proceedings of the Eighth Canadian Conference*, edited by C.P Burgess and R.C. Meyers, (AIP Press, Melville, New York, 1999), pages 301–305. [arXiv:gr-qc/9908023].
- [8] R. Penrose, R. D. Sorkin and E. Woolgar, “A Positive mass theorem based on the focusing and retardation of null geodesics,” arXiv:gr-qc/9301015.
- [9] M. Sereno, “Gravitational lensing by stars with angular momentum,” *Mon. Not. Roy. Astron. Soc.* **344** (2003) 942 [arXiv:astro-ph/0307243].
M. Sereno, “Gravitational lensing in metric theories of gravity,” *Phys. Rev. D* **67** (2003) 064007 [arXiv:astro-ph/0301290].
M. Sereno and V. F. Cardone, “Gravitational lensing by spherically symmetric lenses with angular momentum,” *Astron. Astrophys.* **396** (2002) 393 [arXiv:astro-ph/0209297].
M. Sereno, “Gravitational lensing by spinning and escaping lenses,” arXiv:astro-ph/0209148.
- [10] T. Harko and M. K. Mak, “Anisotropic relativistic stellar models,” *Annalen Phys.* **11** (2002) 3 [arXiv:gr-qc/0302104].
- [11] S. Bharadwaj and S. Kar, “Modelling galaxy halos using dark matter with pressure,” *Phys. Rev. D* **68** (2003) 023516 [arXiv:astro-ph/0304504].
K. Lake, “Galactic Potentials,” *Phys. Rev. Lett.* **92**, 051101 (2004) [arXiv:gr-qc/0302067].